

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 15/82, 15/54, 5/10, A01H 5/00		A1	(11) International Publication Number: WO 96/21738
			(43) International Publication Date: 18 July 1996 (18.07.96)
(21) International Application Number: PCT/US96/00562		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority Data: 08/372,200 15 January 1995 (15.01.95) US		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant: CALGENE, INC. [US/US]; 1920 Fifth Street, Davis, CA 95616 (US).			
(72) Inventor: SHEWMAKER, Christine, K.; 1409 Spring Creek, Woodland, CA 95695 (US).			
(74) Agents: SCHWEDLER, Carl, J. et al.; Calgene, Inc., 1920 Fifth Street, Davis, CA 95616 (US).			
(54) Title: SOLUBLE SOLIDS MODIFICATION USING SUCROSE PHOSPHATE SYNTHASE ENCODING SEQUENCES			
(57) Abstract This invention relates to methods for the utilization of sucrose phosphate synthase encoding sequences to modify the soluble solids in plant sink tissue. The method permits an increase in the sweetness of tomato fruit.			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SOLUBLE SOLIDS MODIFICATION USING
SUCROSE PHOSPHATE SYNTHASE ENCODING SEQUENCES

INTRODUCTION

5

Related Application Data

The present application is a continuation-in-part of U.S. Patent Application Serial No. 08/372,200.

10

Technical Field

The present invention is directed to compositions and methods related to modification of the sucrose biosynthesis pathway to alter the soluble solids content in selected plant tissue.

15

Background

With the development of genetic engineering techniques, it is now possible to transfer genes from a variety of organism into the genome of a large number of different plant species.

20

This process has many advantages over plant breeding techniques, as genes may now be transferred from one plant species to another plant species, rather than simply from a plant to the same, or different, but closely related, species.

25

Sucrose is one of the primary end products of photosynthesis in higher plants. It is also the major carbohydrate transported to sucrose accumulating, or carbon sink, tissues for plant growth and development (Pate, (1976) In: *Transport and Transfer Processes in Plants*, pp. 253-289, Wardlaw, J.F., Passioura, J.B., eds. Academic Press, New York).

SUBSTITUTE SHEET (RULE 26)

Plant regions, such as leaf tissue, where sucrose is synthesized are commonly referred to as sucrose source tissue. Plant storage organs, such as roots or tubers, and fruits are examples of sink tissues.

- 5 Sucrose phosphate synthase (SPS) cDNA sequence, SPS constructs and transgenic SPS tomato lines are described in copending Application, Serial No. 08/175,471. This information is also published in Worrell et al. (*The Plant Cell* (1991) 3:1121-1130), incorporated herein by reference. In particular,
- 10 the copending application describes a maize SPS under the control of the Rubisco small subunit promoter (SSU promoter) from tobacco, providing preferential expression in leaf tissue. Maximum activity of SPS is shown to be significantly increased
- 15 absolute levels of starch and sucrose in the leaves are altered in the predicted manner.

Relevant Literature

- SPS encoding sequences and the generation of transgenic tomato lines from the pCGN3812 SSU-SPS construct is described by
- 20 Worrell et al. (*The Plant Cell* (1991) 3:1121-1130).

Galtier et al. (*Plant Physiol.* (1993) 101:535-543) examined the photosynthetic characteristics of SPS transformants.

- In PCT Application WO 94/00563 antisense potato SPS is placed behind a tuber promoter and used to alter the sucrose
- 25 levels in potato.

Acid invertase encoding sequences are described by Klann et. al., (*Plant Phys.* (1992) 99:351-353.)

SUBSTITUTE SHEET (RULE 26)

SUMMARY OF THE INVENTION

By this invention, a method is disclosed whereby a construct encoding sucrose phosphate synthase (SPS) can be used to modify the solids content of plant sink tissue. Soluble solids include simple sugars, but also can include certain soluble polymers, and other soluble cell components. Total solids includes more complex carbon compounds, such as starches and cellulose. In one embodiment of the present invention, methods are disclosed for increasing the sweetness of fruit.

10 The method provided for increasing the total solids in a plant sink tissue will modify total solids from a given ratio of total solids per unit weight of sink tissue, as measured in control plant cells, to a different ratio of total solids per unit weight of sink tissue. The amount of sucrose available to growing tissues in the plant is increased, and the increased sucrose results in increased total solids per unit weight in the sink tissues of the plant. The method generally comprises growing a plant having integrated into its genome a construct comprising as operably linked components in the 5' to 3' direction of transcription, a transcription initiation region functional in a plant cell and a DNA encoding SPS.

20 The construct transcription initiation region may be constitutive or tissue specific, i.e., preferentially expressed and functional in cells of a particular plant tissue, for instance fruit or leaf. Many such tissue specific promoter regions are known, such as the Rubisco small subunit promoter preferentially expressed in leaf tissue, the patatin promoter expressed in potato tubers and the E8 promoter specific for fruit tissue.

In one embodiment the method produces sink tissue having increased carbon as soluble solids, as an increased ratio of soluble solids per unit weight of sink tissue, as compared to that measured in control plant cells. This results from the increased levels of sucrose generating an increased rate of transportation of the available sucrose into the carbon sink tissue.

In another embodiment a method is provided for modifying the soluble solids ratios in sink tissue, such as the ratio of sucrose to fructose, as compared to that measured in control plant cells or tissue. If the increased soluble solids in said sink tissue comprises fructose, a phenotype will result having an increased sweetness as opposed to the control tissue. Methods are also disclosed, however, whereby a decreased ratio of fructose to sucrose, and whereby a reduced sweetness phenotype may be produced.

The use of constructs comprising encoding sequences to other sucrose metabolizing enzymes, such as acid invertase, or the utilization of such enzymes which are endogenous to the plant sink cells, can be advantageously used with this invention. For instance, acid invertase can be expressed in the cells of sink tissue from an expression construct, or, alternatively, the sink tissue can be prevented from converting sucrose to fructose and glucose by the use of an antisense acid invertase construct, whereby cells of the sink tissue will have a decreased acid invertase activity, and thereby a decreased ratio of fructose to sucrose as compared to cells in a control sink tissue.

Considered part of this invention are the plants, plant cells and plant sink tissue transformed by nucleic acid sequences to SPS demonstrating a modified carbon solids content. The modification can be an increase in one or more solids components, or a change to the ratios of solids components.

In a particular embodiment, the invention provides fruit having increased total soluble solids and/or modified or increased fructose levels, as measured per unit weight.

A preferred embodiment includes fruit having a modified sweetness phenotype, either from a total increase in sweetness by percentage of fruit weight, or from an increased ratio of fructose to sucrose in the soluble solids.

Transformed plant cells of this invention are obtainable through plant transformation techniques which utilize *Agrobacterium* to transfer DNA to the plant cells or alternatively through direct transfer techniques such as electroporation, microinjection or DNA bombardment. In either case, plant carbon sink tissue, such as fruit, may be obtained which has increased or decreased soluble solids content and/or modified sweetness properties.

DETAILED DESCRIPTION OF THE INVENTION

This invention is directed to a method for producing plant cells demonstrating modified soluble solids content. The method utilizes a DNA sequence encoding SPS integrated in the cellular genome as the result of genetic engineering. Plant fruit which contains increased levels of soluble solids and an altered sweetness phenotype is also contemplated in this invention. The

gene encoding SPS is described in copending Application Serial Number 08/175,471, filed on December 27, 1993.

The mechanism whereby the expression of exogenous SPS modifies carbon relationships derives from source-sink relationships. For instance, in those situations where SPS is expressed in leaves, plants have higher sugar and less starch in the leaves. The leaf tissue is a sucrose source, and if this higher sucrose is transported to a sink it results in increased storage carbon (sugars, starch, etc.) per given weight of the sink tissue.

The increased soluble solids in transgenic tomato lines transformed with SPS behind an SSU promoter showed an increase in glucose and fructose over sucrose. Similarly, when the fruit specific E8 promoter was used to express SPS, the resultant increase was in the form of the glucose and fructose.

By this invention, then, the increased carbon produced in a plant sink tissue in the form of soluble solids may be in the form of an increase in glucose and fructose levels as an alternative to, or as well as in, changes to levels of sucrose. In the tomato fruit glucose and fructose are produced from sucrose due to a vacuolar acid invertase that comes on during ripening. Acid invertase converts sucrose to glucose and fructose, and is known to affect fruit sweetness (Hubbard et al. (1991) *Physiol. Plant.* 82:191-196). It was not known prior to the invention whether an increase flow of sucrose into the fruit of tomato would be converted by exogenous acid invertase levels in this manner.

Fructose is twice as sweet, on a molar basis, as glucose, thus the invention provides a mechanism by which to selectively

increase fructose in fruits for increasing sweetness. Thus, in one embodiment, this invention produces tomato fruit (a sink tissue) having increased sweetness.

Thus, the method of altering sink tissue solids with sequences to SPS may be advantageously used in conjunction with endogenous sucrose metabolizing enzymes, whereby sequences to SPS are employed as the sole transformed encoding sequence. The method may also be employed in conjunction with other transformed sequences, for instance sequences encoding other sucrose metabolizing enzymes. Inhibition of certain sucrose metabolizing enzymes could result from the use of antisense expression. The inhibition of acid invertase in tomato fruit, for instance, can lead to fruit having elevated levels of sucrose in the tomato fruit. The sequence to acid invertase is known (Klann et al., (1992) *Plant. Phys.* (1992) 99:351-353). Expression of other sucrose metabolizing enzymes may result in alterations to other carbon components, for instance the expression of starch synthesizing enzymes to act in concert with the increase availability of sucrose may result in increased starch levels in the sink tissue. Such enzymes are known to the art, with many described in copending application having the serial number 08/016,881, the teachings of which are incorporated herein by reference. The transformation of plants using glycogen synthesis enzymes (glgA, glgB and glgC) to modify starch compositions is described in U.S. Patent No. 5,349,123.

A sucrose metabolizing enzyme considered in this invention includes any sequence of amino acids, such as protein, polypeptide, or peptide fragment, which demonstrates the ability to catalyze a reaction involved in the synthesis or degradation

of sucrose or a precursor of sucrose. These can be endogenous plant sequences, by which is meant any sequence which can be naturally found in a plant cell, including native (indigenous) plant sequences as well as sequences from plant viruses or pl
5 pathogenic bacteria, such as *Agrobacterium* or *Rhizobium* species that are naturally found and functional in plant cells.

It will be recognized by one of ordinary skill in the art that sucrose metabolizing enzyme sequences may also be modified using standard techniques of site specific mutation or PCR, or
10 modification of the sequence may be accomplished in producing a synthetic nucleic acid sequence and will still be considered a sucrose biosynthesis enzyme nucleic acid sequence of this invention. For example, wobble positions in codons may be changed such that the nucleic acid sequence encodes the same
15 amino acid sequence, or alternatively, codons can be altered such that conservative amino acid substitutions result. In either case, the peptide or protein maintains the desired enzymatic activity and is thus considered part of the instant invention.

20 A nucleic acid sequence to a sucrose metabolizing enzyme may be a DNA or RNA sequence, derived from genomic DNA, cDNA, mRNA, or may be synthesized in whole or in part. The structural gene sequences may be cloned, for example, by isolating genomic DNA from an appropriate source, and amplifying and cloning the
25 sequence of interest using a polymerase chain reaction (PCR). Alternatively, the gene sequences may be synthesized, either completely or in part, especially where it is desirable to provide plant-preferred sequences. Thus, all or a portion of the desired structural gene may be synthesized using codons

preferred by a selected plant host. Plant-preferred codons may be determined, for example, from the codons used most frequently in the proteins expressed in a particular plant host species.

Other modifications of the gene sequences may result in mutants
5 having slightly altered activity. Once obtained, a sucrose metabolizing enzyme may be utilized with the SPS sequence in a variety of ways.

Other endogenous plant sequences may be useful in nucleic acid constructs of this invention, for example to provide for
10 transcription of the sucrose metabolizing enzyme sequences. Transcriptional regulatory regions are located immediately 5' to the DNA sequences of the gene of interest, and may be obtained from sequences available in the literature, or identified and characterized by isolating genes having a desirable
15 transcription pattern in plants, and studying the 5' nucleic acid sequences. Numerous transcription initiation regions which provide for a variety of constitutive or regulatable, e.g. inducible, expression in a plant cell are known. Among sequences known to be useful in providing for constitutive gene
20 expression are regulatory regions associated with *Agrobacterium* genes, such as for nopaline synthase (*Nos*), mannopine synthase (*Mas*), or octopine synthase (*Ocs*), as well as regions coding for expression of viral genes, such as the 35S and 19S regions of cauliflower mosaic virus (*CaMV*). The term constitutive as used
25 herein does not necessarily indicate that a gene is expressed at the same level in all cell types, but that the gene is expressed in a wide range of cell types, although some variation in abundance is often detectable.

In providing for transcription and/or expression of the sucrose metabolizing enzyme sequences, for various reasons one may wish to limit the expression of these enzymes to plant cells which function as carbon sinks. Towards this end, one can

5 identify useful transcriptional initiation regions that provide for expression preferentially in specific tissue types, such as roots, tubers, seeds or fruit. These sequences may be identified from cDNA libraries using differential screening techniques, for example, or may be derived from sequences known

10 in the literature. Useful transcriptional initiation regions preferentially provide for transcription in certain tissues or under certain growth conditions, such as those from napin, seed or leaf ACP, the small subunit of RUBISCO, patatin, zein, and the like. Fruit specific promoters are also known, one such

15 promoter is the E8 promoter, described in Deikman et al. (1988) *EMBO J.* 2:3315-3320; and DellaPenna et al. (1989) *Plant Cell* 1:53-63, the teachings of which are incorporated herein by reference. An E8-SPS construct (fruit-specific promoter) will express SPS in a fruit-specific manner, whereby the levels of

20 sucrose produced in the fruit may be elevated. If coupled with antisense invertase, the increase in sucrose would be maintained. This is a particular issue in tomatoes where acid invertase present in the fruit drives the production of glucose and fructose from sucrose.

25 Sequences to be transcribed are located 3' to the plant transcription initiation region and may be oriented, in the 5'-3' direction, in the sense orientation or the antisense orientation. In the sense orientation, an mRNA strand is produced which encodes the desired sucrose metabolizing enzyme.

while in antisense constructs, an RNA sequence complementary to an enzyme coding sequence is produced. The sense orientation is desirable when one wishes to produce the sucrose metabolizing enzyme in plant cells, whereas the antisense strand may be
5 useful to inhibit production of a related plant sucrose metabolizing enzymes. The presence of sucrose metabolizing enzyme sequences in the genome of the plant host cell may be confirmed, for example by a Southern analysis of DNA or a Northern analysis of RNA sequences or by PCR methods.

10 In addition to sequences providing for transcriptional initiation in a plant cell, also of interest are sequences which provide for transcriptional and translational initiation of a desired sequence encoding a sucrose metabolizing enzyme. Translational initiation regions may be provided from the source
15 of the transcriptional initiation region or from the gene of interest. In this manner, expression of the sucrose metabolizing enzyme in a plant cell is provided. The presence of the sucrose metabolizing enzyme in the plant host cell may be confirmed by a variety of methods including a immunological
20 analysis of the protein (e.g. Western or ELIZA), as a result of phenotypic changes observed in the cell, such as altered soluble solids content or by assay for increased enzyme activity, and the like. Other sequences may be included in the nucleic acid construct providing for expression of the sucrose metabolizing
25 enzymes ("expression constructs") of this invention, including endogenous plant transcription termination regions which will be located 3' to the desired sucrose metabolizing enzyme encoding sequence. For instance, transcription termination sequences derived from a patatin gene may be utilized when the sink tissue

is potato tubers. Transcription termination regions may also be derived from genes other than those used to regulate the transcription in the nucleic acid constructs of this invention. Transcription termination regions may be derived from a variety of different gene sequences, including the *Agrobacterium*, viral and plant genes discussed above for their desirable 5' regulatory sequences.

Further constructs are considered which provide for transcription and/or expression of more than one sucrose metabolizing enzyme. For example, one may wish to provide enzymes to plant cells of the sink tissue which provide for modification of the type of soluble solids to be produced therein, as well as for enhancing or otherwise modifying the increase or decrease in overall soluble solids production. An example of enzymes which may prove useful in modifying soluble solids ratios is the acid invertase enzyme.

In developing the nucleic acid constructs of this invention, the various components of the construct or fragments thereof will normally be inserted into a convenient cloning vector, e.g. a plasmid, which is capable of replication in a bacterial host, e.g. *E. coli*. Numerous vectors exist that have been described in the literature, many of which are commercially available. After each cloning, the cloning vector with the desired insert may be isolated and subjected to further manipulation, such as restriction, insertion of new fragments or nucleotides, ligation, deletion, mutation, resection, etc. so as to tailor the components of the desired sequence. Once the construct has been completed, it may then be transferred to an

appropriate vector for further manipulation in accordance with the manner of transformation of the host cell.

The constructs of this invention providing for transcription and/or expression of sucrose metabolizing enzyme sequences of this invention may be utilized as vectors for plant cell transformation. The manner in which nucleic acid sequences are introduced into the plant host cell is not critical to this invention. Direct DNA transfer techniques, such as electroporation, microinjection or DNA bombardment may be useful. To aid in identification of transformed plant cells, the constructs of this invention may be further manipulated to include plant selectable markers. The use of plant selectable markers is preferred in this invention as the amount of experimentation required to detect plant cells is greatly reduced when a selectable marker is expressed. Useful selectable markers include enzymes which provide for resistance to an antibiotic such as gentamycin, hygromycin, kanamycin, and the like. Similarly, enzymes providing for production of a compound identifiable by color change, such as GUS, or luminescence, such as luciferase are useful.

An alternative method of plant cell transformation employs plant vectors which contain additional sequences which provide for transfer of the desired sucrose metabolizing enzyme sequences to a plant host cell and stable integration of these sequences into the genome of the desired plant host. Selectable markers may also be useful in these nucleic acid constructs to provide for differentiation of plant cells containing the desired sequences from those which have only the native genetic material. Sequences useful in providing for transfer of nucleic

acid sequences to host plant cells may be derived from plant pathogenic bacteria, such as *Agrobacterium* or *Rhizogenes*, plant pathogenic viruses, or plant transposable elements.

When *Agrobacterium* is utilized for plant transformation, 5 may be desirable to have the desired nucleic acid sequences bordered on one or both ends by T-DNA, in particular the left and right border regions, and more particularly, at least the right border region. These border regions may also be useful when other methods of transformation are employed.

10 Where *Agrobacterium* or *Rhizogenes* sequences are utilized for plant transformation, a vector may be used which may be introduced into an *Agrobacterium* host for homologous recombination with the T-DNA on the Ti- or Ri-plasmid present in the host. The Ti- or Ri- containing the T-DNA for recombination 15 may be armed (capable of causing gall formation), or disarmed (incapable of causing gall formation), the latter being permissible so long as a functional complement of the vir gene which encode trans-acting factors necessary for transfer of DNA to plant host cells, is present in the transformed *Agrobacterium* 20 host. Using an armed *Agrobacterium* strain can result in a mixture of normal plant cells, some of which contain the desired nucleic acid sequences, and plant cells capable of gall formation due to the presence of tumor formation genes. Cells containing the desired nucleic acid sequences, but lacking tumor 25 genes can be selected from the mixture such that normal transgenic plants may be obtained.

In a preferred method where *Agrobacterium* is used as the vehicle for transforming host plant cells, the expression or transcription construct bordered by the T-DNA border region(s) 14

will be inserted into a broad host range vector capable of replication in *E. coli* and *Agrobacterium*, there being broad host range vectors described in the literature. Commonly used is pRK2 or derivatives thereof. See, for example, Ditta, et al., 5 (Proc. Nat. Acad. Sci., U.S.A. (1980) 77:7347-7351) and EPA 0 120 515, which are incorporated herein by reference.

Alternatively, one may insert the sequences to be expressed in plant cells into a vector containing separate replication sequences, one of which stabilizes the vector in *E. coli*, and 10 the other in *Agrobacterium*. See, for example, McBride and Summerfelt (Plant Mol. Biol. (1990) 14:269-276), wherein the pRIHRI (Jouanin, et al., Mol. Gen. Genet. (1985) 201:370-374) origin of replication is utilized and provides for added stability of the plant expression vectors in host *Agrobacterium* 15 cells.

Utilizing vectors such as those described above, which can replicate in *Agrobacterium* is preferred. In this manner, recombination of plasmids is not required and the host *Agrobacterium* vir regions can supply trans-acting factors 20 required for transfer of the T-DNA bordered sequences to plant host cells.

In general, the plant vectors of this invention will contain sucrose metabolizing enzyme sequence(s) and sequences providing for transcription or expression of these sequences in 25 a plant host cell. The plant vectors containing the desired sequences may be employed with a wide variety of plant cells and plants. Fruit producing plants and plants which produce and store reserve starch are of particular interest, the latter including, but in no way limited to, corn, cereal grains,

sorghum, rice, potato, tapioca, cassava, arrowroot and sago. The method may also be useful in increasing the sweetness in fruits, including, but in no way limited to, tomato, strawberry and melon.

5 Also considered part of this invention are plants containing the nucleic acid sequences of this invention, and following from that, plants modified by the expression of SPS and possibly additional sucrose metabolizing enzymes as the result of expression of the sequences in plant cells, or having
10 a decreased expression of a such by the inhibition of such additional sucrose metabolizing enzymes from antisense expression of the encoding sequences. Methods of regenerating whole plants from plant cells are known in the art, and the method of obtaining transformed and regenerated plants is not
15 critical to this invention. In general, transformed plant cells are cultured in an appropriate medium, which may contain selective agents such as antibiotics, where selectable marker are used to facilitate identification of transformed plant cells. Once callus forms, shoot formation can be encouraged by
20 employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be used to establish repetitive generations either from seed or using vegetative propagation techniques.

25 The invention now being generally described, it will be more readily understood by reference to the following examples which are included for purposes of illustration only and are not intended to limit the present invention.

Measurement of specific gravity or free sugar content may be useful to detect modified sweetness or total soluble solids, with other methods, such as HPLC and gel filtration, also being useful.

5

EXAMPLES

Constructs and plants with the designations 3812 and 3815 are as described in copending Application Serial No. 08/175,471, which in turn is a continuation of U.S. Patent Application
10 Serial No. 07/672,646, both of which are incorporated by reference. Briefly, the construct pCGN3812 contains the construct 5'-35S-nptII-tml-3'/5'=tobacco Rubisco SSU-maize SPS cDNA-3'. The construct pCGN3815 contains the construct 5'-35S-nptII-tml-3'/5'35S-maize SPS cDNA-3'. Tomato lines arising from
15 separate transformation events are signified by a hyphen and a number following the construct designation.

Example 1 - Soluble Solids in T2 SSU-SPS Plants

Investigation of the soluble solids in the fruits of the
20 SSU-SPS lines was initially done on extracts from fruit of 3812-9 and 3812-11 lines grown in a Biotron incubator. T2 plants were illuminated by metal halide lamps at a peak level of 500 μmol photons/m/s (pot level), 26 C for the 16 h day and 18 C at night, and a relative humidity of 60 %. Plants were watered
25 daily with half-strength Hoagland's solution (Hoagland and Arnon, Calif. Agricult. Exp. Sta. Cir. (1938) 357:1-39). These lines were segregating as the original lines contained at least 2 insertions.

Brix analysis (soluble solids) on extracts from these plants revealed lines with Brix readings as much as 40% higher than the controls. The extracts measured were the average of 3 fruit from one plant.

- 5 Measurements were also taken for fruit from a segregating T2 population of 3812-11 plants in the greenhouse. The controls averaged a Brix reading of 3.5 while the transgenics averaged 4.0, an increase of 14%.

10 **Example 2 - Homozygous Plants**

- T4 homozygous lines were generated from original 3812-9 transformants in UC82-B tomatoes. The original line segregated 15:1 for Kan resistance, indicating that it had two insertion sites. Two homozygous lines were generated and verified to be
15 different by Southern border analysis. These lines were designated A and B.

- Individual homozygote (T4) lines were grown in the greenhouse, with three fruit taken from each plant and 3 plants analyzed from each line. The Brix of the UC82B controls was
20 3.35 while the Brix on the 3812-9 lines ranged from 3.7 to 4.1. This is an increase from 12 % to 24 %. Statistics (LSD) on all the lines in which fruit from 3 plants were analyzed showed these results to be significant at a .01% level (99%).

- Measurements were also made on homozygous lines of tomato
25 plants transformed with the 35S CaMV promoter-SPS construct pCGN3815. In one line, 3815-13-2 there was a substantial increase in yield of tomatoes, in terms of an increase in both fruit size and in fruit number, as measured against non-transformed control plants and as against SSU-SPS homozygous

line controls. The 3815-13-2 plants also produced a second flush of fruit. A second transgenic line containing the pCGN3815 construct did not produce these dramatic yield increases.

5

Example 3 - Brix Analysis of Field Trial SSU-SPS Fruit

Field trial results of RI measurements are provided in Table 1. The R/I (refractive index) was measured several times on the fruit of these plants (Table 1). R/I is a measure of soluble solids and is indicative of sugars and acids. The transgenic A lines consistently had a higher R/I (3.9 to 4.1) than the control. (3.5).

10

Example 4 - HPLC Analysis on SSU-SPS Fruit Sugars

Fruit from the SPS plants described in Example 3 were further analyzed by HPLC to determine contributions of individual sugars to the increased soluble solids content. As seen in Table 2, sucrose did not increase as much as might be expected based on the fact that sucrose is the sugar transported by the plant into the fruit. Glucose was not increased as much as fructose, which increased nearly 50 %.

It is evident from the above results, that plant cells and plants can be produced which have improved properties or may produce a desired phenotype. In accordance with the subject invention, it is now seen that SPS sequences may be introduced into a plant host cell and be used to express the enzyme to increase soluble solids content in fruit. Moreover, it is seen that the SPS may be used to alter the overall content and ratio of soluble solids in plant sink tissue, resulting in a demonstrable phenotype in planta, such as altered fruit sweetness. In this manner, fruits, such as tomato fruit, having modified sweetness may be obtained.

Example 5 - Fruit Specific Expression of SPS

E8-SPS constructs designated as pCGN3342 and pCGN3343 contain the tomato E8 promoter comprising the approximately 2.1 kb 5' region of the E8 promoter. A description of this promoter region can be found in Deikman et al., *supra*, and in Deikman et al. (Plant Physiol. (1992) 100:2013-2017).

This E8 promoter is fused to the same SPS encoding sequence used for pCGN3812 and pCGN3815, only the SPS sequence used in these constructs has been truncated at the ApoI site just 3' of

the SPS encoding sequence (at nucleotide 3318), and fused to a 1.2 kb region of the *tml* 3' region from pTiA6 (Barker et al., (1983) Plant Mol. Biol. 2:335-350; sequence 11207-10069 of the T-DNA region from the *Agrobacterium tumefaciens* Ti plasmid pTi15955). Constructs pCGN3342 and pCGN3343 are the opposite orientations of this E8-SPS-*tml* construct in the 35S kan binary, pCGN1557 (McBride and Summerfelt, supra). Tomato lines arising from separate transformation events using pCGN3342 and pCGN3343 are signified by the construct number followed by a hyphen and an event number.

Table 3 provides data from RI measurements of soluble solids in tomatoes from greenhouse studies of T1 plants. The R/I (refractive index) was measured several times on the fruit of these plants.

Assays were made for the SPS activity in control and transgenic fruit from the 3343-6 and 3342-11 events. The control 7060 fruit was assayed with a SPS activity rate of 17.8 μ moles sucrose/gram fresh weight/hour. Activity was much higher in the transgenic lines, with the 3343-6 event having a rate of 67.5 μ moles sucrose/g/hr and 3342-11 event measured at 36.6 μ moles sucrose/g/hr. This demonstrates that the increase sugar concentration of fruit in transgenic tomatoes over the control correlates to an increase of SPS activity in the fruit.

Tables 4 and 5 provide an analysis of individual sugars as measured by HPLC from two separate trials, to determine contributions of each sugar to the increased soluble solids content observed in transgenic E8-SPS fruit. The data of table 4 and 5 demonstrate that increased SPS activity from transgenic expression in fruit by a fruit specific promoter can produce an

overall net increase in sugars in the fruit. Due to the endogenous acid invertase found in ripening tomato fruit, increases in sugar are found in glucose and fructose.

It also appears that there is a correlating increase in acid levels with an increase in sugar content in fruit transformed with E8-SPS.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference:

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be readily apparent to those of ordinary skill in the art in light of the teaching of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

CLAIMS

5

What is claimed is:

1. A method of increasing the total solids in a plant sink tissue from a given ratio of total solids per unit weight of sink tissue, as measured in control plant cells, to a different
10 ratio of total solids per unit weight, wherein said method comprises:

growing a plant having integrated into its genome a construct comprising, as operably linked components in the 5' to 3' direction of transcription, a transcription initiation region
15 functional in a plant cell and a DNA encoding a sucrose phosphate synthase, wherein said DNA encoding a sucrose phosphate synthase is not naturally linked to said transcription initiation region, and wherein said cell is grown under conditions which will permit said transcription initiation
20 region to function,

whereby the amount of sucrose available to growing tissues in said plant is increased, and whereby the increased sucrose results in increased total solids per unit weight in said sink tissue.

25 2. The method of Claim 1 wherein said DNA encoding a sucrose phosphate synthase encodes a sucrose phosphate synthase derived from corn.

3. The method of Claim 1 wherein said transcription initiation region is tissue specific.

4. The method of Claim 3 wherein said transcription initiation region is functional in a fruit cell.

5. The method of Claim 3 wherein said transcription initiation region is functional in a leaf cell.

5 6. The method of Claim 5 wherein said transcription initiation region is a Rubisco small subunit promoter.

7. The method of Claim 1 wherein said increased solids are soluble solids.

8. The method of Claim 1 wherein said tissue is a fruit.

10 9. The method of Claim 8 wherein said plant is a tomato plant and wherein said plant sink tissue is a tomato fruit.

10. A method of increasing the soluble solids in a plant sink tissue from a given ratio of soluble solids per unit weight of sink tissue, as measured in control plant cells, to a
15 different ratio of soluble solids per unit weight, wherein said method comprises:

growing a plant having integrated into its genome a construct comprising, as operably linked components in the 5' to 3' direction of transcription, a transcription initiation region
20 functional in a plant cell and a DNA encoding a sucrose phosphate synthase, wherein said DNA encoding a sucrose phosphate synthase is not naturally linked to said transcription initiation region, and wherein said cell is grown under conditions which will permit said transcription initiation
25 region to function,

whereby starch is converted to sucrose in said cell and whereby an increased level of sucrose is made available to said plant sink tissue.

11. The method of Claim 10 wherein said DNA encoding a sucrose phosphate synthase encodes a sucrose phosphate synthase derived from corn.

12. The method of Claim 10 wherein said transcription
5 initiation region is tissue specific.

13. The method of Claim 12 wherein said transcription initiation region is functional in a fruit cell.

14. The method of Claim 12 wherein said transcription initiation region is functional in a leaf cell and wherein said
10 sucrose is transported into said sink tissue.

15. The method of Claim 14 wherein said transcription initiation region is a Rubisco small subunit promoter.

16. The method of Claim 10 wherein said tissue is a fruit.

17. The method of Claim 16 wherein said plant is a tomato
15 plant and wherein said plant sink tissue is a tomato fruit.

18. A tomato fruit produced by the method of Claim 17.

19. The method of Claim 11 wherein said increased soluble solids component in said sink tissue comprises sucrose.

20. The method of Claim 11 wherein said increased soluble
20 solids in said sink tissue comprises glucose and fructose.

21. A method of modifying the soluble solids ratios in a cell of a plant sink tissue, as compared to that measured in control plant cells, to a different ratio of soluble solids, wherein said method comprises:

25 growing a plant having a sucrose metabolizing enzyme in cells of said sink tissue capable of acting on sucrose to alter sucrose levels in said sink tissue, said plant having integrated into its genome a construct comprising, as operably linked components in the 5' to 3' direction of transcription, a

transcription initiation region functional in a plant cell and a DNA encoding a sucrose phosphate synthase, wherein said DNA encoding a sucrose phosphate synthase is not naturally linked to said transcription initiation region, and wherein said cell is
5 grown under conditions which will permit said transcription initiation region to function,

whereby starch is converted to sucrose in said cell to provide an increased level of sucrose to said plant sink tissue and whereby said sucrose is acted on by said sucrose
10 metabolizing enzyme.

22. The method of Claim 21 wherein cells of said sink tissue have an increased ratio of fructose to sucrose as compared to that of a cell in a control sink tissue.

23. The method of Claim 21 wherein said transcription
15 initiation region is tissue specific.

24. The method of Claim 21 wherein said transcription initiation region is functional in a fruit cell.

25. The method of Claim 21 wherein said transcription initiation region is functional in a leaf cell and wherein said
20 sucrose is transported into said sink tissue.

26 The method of Claim 25 wherein said transcription initiation region is a Rubisco small subunit promoter.

27. A plant cell having modified levels of fructose and sucrose, wherein said modified levels of fructose and sucrose
25 are produced according to the method of Claim 21.

28. The method of Claim 21 wherein cells of said sink tissue have a decreased ratio of fructose to sucrose as compared to that of a cell in a control sink tissue.

29. The method of Claim 21 whereby said sucrose metabolizing enzyme is acid invertase present in cells of said sink tissue, whereby sucrose is degraded to fructose and glucose.

5 30. The method of Claim 29 wherein said acid invertase is endogenous to said cells of said sink tissue.

31. The method of Claim 29 wherein said acid invertase is expressed in said cells of said sink tissue from an expression construct, wherein said construct comprises, as operably linked
10 components in the 5' to 3' direction of transcription, a transcription initiation region functional in said sink tissue cell and a DNA encoding a acid invertase, wherein said cell is grown under conditions which will permit said transcription initiation region to function.

15 32. The method of Claim 21 wherein said method further comprises;

) growing said plant having integrated into its genome a second construct comprising, as operably linked components in the 5' to 3' direction of transcription, a transcription
20 initiation region functional in said sink tissue cell an acid invertase gene sequence, wherein said cell is grown under conditions which will permit said transcription initiation region to function, and wherein said acid invertase gene sequence is in an antisense orientation with respect to said
25 transcription initiation region, '

whereby said sink tissue is prevented from converting sucrose to fructose and glucose and whereby cells of said sink tissue have a decreased ratio of fructose to sucrose as compared to that of cells in a control sink tissue.

33. The method of Claim 32 wherein said transcription initiation region linked to said antisense DNA sequence to acid invertase is a double 35S promoter.

34. A method of increasing the sweetness of fruit in a transgenic plant, as compared to that measured in tissue of fruit of control plants, said method comprising:

growing a plant having integrated into its genome a construct comprising, as operably linked components in the 5' to 3' direction of transcription, a fruit specific transcription initiation region and a DNA sequence encoding sucrose phosphate synthase, wherein said sucrose phosphate synthase encoding sequence is not naturally linked to said fruit specific transcription initiation region, and wherein said cell is grown under conditions which will permit said fruit specific transcription initiation region to function and increase the overall sugar content in cells of said fruit.

35. The method of Claim 1 wherein cells of said fruit have an increased level of sucrose as compared to that of a cell in a control fruit.

36. The method of Claim 1 wherein cells of said fruit have an increased level of fructose as compared to that of a cell in a control fruit.

37. The method of Claim 1 wherein cells of said fruit have an increased level of glucose as compared to that of a cell in a control fruit.

38. The method of Claim 1 wherein said fruit specific transcription initiation region is from the E8 promoter.

39. The method of Claim 1 wherein said plant is a tomato.

40. A plant fruit cell having increased sugar content produced according to the method of Claim 1.

41. A tomato fruit cell according to Claim 7.

42. A plant comprising a cell according to Claim 7.

5 43. A tomato plant according to Claim 9.

44. The method of Claim 11 wherein said method further comprises;

growing said plant having integrated into its genome a second construct comprising, as operably linked components in
10 the 5' to 3' direction of transcription, a transcription initiation region functional in a plant fruit cell and a DNA sequence encoding acid invertase, wherein said cell is grown under conditions which will permit said transcription initiation region of said second construct to function and express acid
15 invertase in fruit of said plant.

45. The method of Claim 11 wherein said transcription initiation region linked to said acid invertase is a fruit specific promoter.

46. The method of Claim 12 wherein said fruit specific
20 promoter is E8.

47. The method of Claim 11 wherein said method further comprises;

growing said plant having integrated into its genome a second construct comprising, as operably linked components in
25 the 5' to 3' direction of transcription, a transcription initiation region functional in a plant fruit cell and a DNA sequence encoding acid invertase, wherein said cell is grown under conditions which will permit said transcription initiation region to function, and wherein said acid invertase gene

sequence is in an antisense orientation with respect to said transcription initiation region,

whereby cells of said plant fruit are prevented from expressing acid invertase and said overall sugar content in
5 cells of said fruit is increased by an increase in sucrose.

48. The method of Claim 14 wherein said transcription initiation region linked to said antisense DNA sequence to acid invertase is a fruit specific promoter.

49. The method of Claim 14 wherein said fruit specific
10 promoter is E8.

50. The method of Claim 14 wherein said transcription initiation region linked to said antisense DNA sequence to acid invertase is a double 35S promoter.

15

INTERNATIONAL SEARCH REPORT

International Application No.
 PCT/US 96/00562

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C12N15/82 C12N15/54 C12N5/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PLANT PHYSIOLOGY, vol. 101, 1993, pages 535-543, XP002002229 GALTIER, N., ET AL.: "Effects of elevated sucrose-phosphate synthase activity on photosynthesis, assimilate partitioning, and growth in tomato (Lycopersicon esculentum var UC82B)" see the whole document</p> <p style="text-align: center;">--- -/--</p>	1-50

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

6 May 1996

Date of mailing of the international search report

10.05.1996

Name and mailing address of the ISA

European Patent Office, P.O. Box 5318 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 631 epo nl,
Fax (+31-70) 340-2016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/US 96/00562

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ANNUAL MEETING OF THE AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, PORTLAND, OREGON, USA, JULY 30-AUGUST 3, 1994. PLANT PHYSIOLOGY (ROCKVILLE) 105 (1 SUPPL.). 1994. 23. , XP002002230 FOYER C H., ET AL.: "Modifications in carbon assimilation, carbon partitioning and total biomass as a result of over-expression of sucrose phosphate synthase in transgenic tomato plants." see abstract 65 ---	1-50
A	WO,A,92 16631 (ROUSSEL UCLAF) 1 October 1992 see the whole document ---	1-50
A	WO,A,94 00563 (INST GENBIOLOGISCHE FORSCHUNG ;SONNEWALD UWE (DE)) 6 January 1994 see the whole document ---	1-50
A	EP,A,0 530 978 (CAMBRIDGE ADVANCED TECH) 10 March 1993 see the whole document ---	1-50
A	EP,A,0 438 904 (CAMBRIDGE ADVANCED TECH) 31 July 1991 see page 3, line 14 - line 20 ---	1-50
A	WO,A,94 28146 (HOECHST SCHERING AGREVO GMBH ;HESSE HOLGER (DE); MUELLER ROEBER BE) 8 December 1994 see the whole document ---	1-50
A	BIOLOGICAL ABSTRACTS, vol. 100, 1995 Philadelphia, PA, US; abstract no. 42942, MICALLEF, B.J., ET AL.: "Altered photosynthesis, flowering, and fruiting in transgenic tomato plants that have an increased capacity for sucrose synthesis" XP002002232 see abstract & PLANTA (HEIDELBERG), vol. 196, no. 2, 1995, pages 327-334, ---	1-50
A	WO,A,93 06711 (UNIV CALIFORNIA) 15 April 1993 see the whole document ---	31,32, 38,44-49

1

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/00562

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,92 14831 (SALK INST BIOTECH IND) 3 September 1992 see page 4, line 1 - page 6, line 21; claim 22 see page 9, line 10 - line 18 see page 59, line 17 - page 60, line 15 ---	31,32, 44-49
A	PLANT PHYSIOLOGY, vol. 100, 1992, pages 310-308, XP002002231 HEINEKE, D., ET AL.: "Apoplastic expression of yeast-derived invertase in potato" see page 307, right-hand column ---	31
A	WO,A,93 14212 (ZENECA LTD) 22 July 1993 see the whole document -----	18,40-43

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Patent Application No.

PCr/US 96/00562

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9216631	01-10-92	AU-B- 662111	24-08-95
		AU-B- 8765591	21-10-92
		EP-A- 0533849	31-03-93
WO-A-9400563	06-01-94	DE-A- 4220758	05-01-94
		AU-B- 4500493	24-01-94
		CA-A- 2136828	06-01-94
		EP-A- 0648262	19-04-95
		HU-A- 71573	28-12-95
		JP-T- 7508406	21-09-95
EP-A-0530978	10-03-93	CA-A- 2075388	09-02-93
EP-A-0438904	31-07-91	AU-B- 6836590	04-07-91
		JP-A- 4341126	27-11-92
		US-A- 5387756	07-02-95
WO-A-9428146	08-12-94	DE-A- 4317596	01-12-94
		EP-A- 0701617	20-03-96
WO-A-9306711	15-04-93	AU-B- 2772792	03-05-93
WO-A-9214831	03-09-92	AU-B- 1456292	15-09-92
		BR-A- 9205480	01-03-94
		EP-A- 0573566	15-12-93
		HU-A- 66831	30-01-95
WO-A-9314212	22-07-93	AU-B- 3262293	03-08-93

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

BLANK PAGE